

Set-theoretic remarks on a possible definition of elementary ∞ -topos

Giulio Lo Monaco

Masaryk University

HoTT, 2019
Pittsburgh, Pennsylvania

16 August 2019

Definition

An ∞ -category \mathcal{X} is called a *geometric ∞ -topos* if there is a small ∞ -category \mathcal{C} and an adjunction

$$\mathcal{P}(\mathcal{C}) \begin{array}{c} \xrightarrow{L} \\ \perp \\ \xleftarrow{i} \end{array} \mathcal{X}$$

where i is full and faithful, $L \circ i$ is accessible and L preserves all finite limits.

Definition

An ∞ -category \mathcal{X} is called a *geometric ∞ -topos* if there is a small ∞ -category \mathcal{C} and an adjunction

$$\mathcal{P}(\mathcal{C}) \begin{array}{c} \xrightarrow{L} \\ \perp \\ \xleftarrow{i} \end{array} \mathcal{X}$$

where i is full and faithful, $L \circ i$ is accessible and L preserves all finite limits.

In particular, every geometric ∞ -topos is presentable.

Ingredients: Dependent sums and products

Let $f : X \rightarrow Y$ a morphism in an ∞ -category \mathcal{E} with pullbacks.

Ingredients: Dependent sums and products

Let $f : X \rightarrow Y$ a morphism in an ∞ -category \mathcal{E} with pullbacks.

- A dependent sum along f is a left adjoint of the base change $f^* : \mathcal{E}_{/Y} \rightarrow \mathcal{E}_{/X}$.

Ingredients: Dependent sums and products

Let $f : X \rightarrow Y$ a morphism in an ∞ -category \mathcal{E} with pullbacks.

- A dependent sum along f is a left adjoint of the base change $f^* : \mathcal{E}_{/Y} \rightarrow \mathcal{E}_{/X}$.
- A dependent product along f , if it exists, is a right adjoint to the base change $f^* : \mathcal{E}_{/Y} \rightarrow \mathcal{E}_{/X}$.

Ingredients: Dependent sums and products

Let $f : X \rightarrow Y$ a morphism in an ∞ -category \mathcal{E} with pullbacks.

- A dependent sum along f is a left adjoint of the base change $f^* : \mathcal{E}_{/Y} \rightarrow \mathcal{E}_{/X}$.
- A dependent product along f , if it exists, is a right adjoint to the base change $f^* : \mathcal{E}_{/Y} \rightarrow \mathcal{E}_{/X}$.

$$\mathcal{E}_{/X} \begin{array}{c} \xrightarrow{\Sigma_f} \\ \perp \\ \xleftarrow{f^*} \\ \perp \\ \xrightarrow{\Pi_f} \end{array} \mathcal{E}_{/Y} : f^*$$

Ingredients: Dependent sums and products

Let $f : X \rightarrow Y$ a morphism in an ∞ -category \mathcal{E} with pullbacks.

- A dependent sum along f is a left adjoint of the base change $f^* : \mathcal{E}/_Y \rightarrow \mathcal{E}/_X$.
- A dependent product along f , if it exists, is a right adjoint to the base change $f^* : \mathcal{E}/_Y \rightarrow \mathcal{E}/_X$.

$$\mathcal{E}/_X \begin{array}{c} \xrightarrow{\Sigma_f} \\ \perp \\ \xleftarrow{f^*} \\ \perp \\ \xrightarrow{\Pi_f} \end{array} \mathcal{E}/_Y : f^*$$

Remark

Dependent sums always exist by universal property of pullbacks.

Ingredients: Dependent sums and products

Let $f : X \rightarrow Y$ a morphism in an ∞ -category \mathcal{E} with pullbacks.

- A dependent sum along f is a left adjoint of the base change $f^* : \mathcal{E}_{/Y} \rightarrow \mathcal{E}_{/X}$.
- A dependent product along f , if it exists, is a right adjoint to the base change $f^* : \mathcal{E}_{/Y} \rightarrow \mathcal{E}_{/X}$.

$$\mathcal{E}_{/X} \begin{array}{c} \xrightarrow{\Sigma_f} \\ \perp \\ \xleftarrow{f^*} \\ \perp \\ \xrightarrow{\Pi_f} \end{array} \mathcal{E}_{/Y} : f^*$$

Remark

Dependent sums always exist by universal property of pullbacks.

Proposition

In a geometric ∞ -topos all dependent products exist.

Let S be a class of morphisms in an ∞ -category \mathcal{E} , which is closed under pullbacks.

A classifier for the class S is a morphism $t : \bar{U} \rightarrow U$ such that for every object X the operation of pulling back defines an equivalence of ∞ -groupoids

$$\mathrm{Map}(X, U) \simeq (\mathcal{E}_{/X}^S)^\sim$$

Definition (Shulman)

An elementary ∞ -topos is an ∞ -category \mathcal{E} such that

Definition (Shulman)

An elementary ∞ -topos is an ∞ -category \mathcal{E} such that

- 1 \mathcal{E} has all finite limits and colimits.
- 2 \mathcal{E} is locally Cartesian closed.
- 3 The class of all monomorphisms in \mathcal{E} admits a classifier.
- 4

Definition (Shulman)

An elementary ∞ -topos is an ∞ -category \mathcal{E} such that

- 1 \mathcal{E} has all finite limits and colimits.
- 2 \mathcal{E} is locally Cartesian closed.
- 3 The class of all monomorphisms in \mathcal{E} admits a classifier.
- 4 For each morphism f in \mathcal{E} there is a class of morphisms $S \ni f$ such that S has a classifier and is closed under finite limits and colimits taken in overcategories and under dependent sums and products.

Definition (Shulman)

An elementary ∞ -topos is an ∞ -category \mathcal{E} such that

- 1 \mathcal{E} has all finite limits and colimits.
- 2 \mathcal{E} is locally Cartesian closed.
- 3 The class of all monomorphisms in \mathcal{E} admits a classifier.
- 4 For each morphism f in \mathcal{E} there is a class of morphisms $S \ni f$ such that S has a classifier and is closed under finite limits and colimits taken in overcategories and under dependent sums and products.

We will only focus on a subaxiom of (4):

Definition

We say that a class of morphisms S satisfies (DepProd) if it has a classifier and it is closed under dependent products

Theorem (Adámek, Rosický for the 1-dimensional case)

Given a small family $(f_i : \mathcal{K}_i \rightarrow \mathcal{L}_i)_{i \in I}$ of accessible functors between presentable ∞ -categories, there are arbitrarily large cardinals κ such that all functors f_i 's preserve κ -compact objects.

Theorem (Adámek, Rosický for the 1-dimensional case)

Given a small family $(f_i : \mathcal{K}_i \rightarrow \mathcal{L}_i)_{i \in I}$ of accessible functors between presentable ∞ -categories, there are arbitrarily large cardinals κ such that all functors f_i 's preserve κ -compact objects.

Example

- *We may assume that κ -compact objects in a presheaf ∞ -category are precisely the objectwise κ -compact presheaves.*

Theorem (Adámek, Rosický for the 1-dimensional case)

Given a small family $(f_i : \mathcal{K}_i \rightarrow \mathcal{L}_i)_{i \in I}$ of accessible functors between presentable ∞ -categories, there are arbitrarily large cardinals κ such that all functors f_i 's preserve κ -compact objects.

Example

- *We may assume that κ -compact objects in a presheaf ∞ -category are precisely the objectwise κ -compact presheaves.*
- *Given a diagram shape R , we may assume that κ -compact objects are stable under R -limits.*

Theorem (Adámek, Rosický for the 1-dimensional case)

Given a small family $(f_i : \mathcal{K}_i \rightarrow \mathcal{L}_i)_{i \in I}$ of accessible functors between presentable ∞ -categories, there are arbitrarily large cardinals κ such that all functors f_i 's preserve κ -compact objects.

Example

- *We may assume that κ -compact objects in a presheaf ∞ -category are precisely the objectwise κ -compact presheaves.*
- *Given a diagram shape R , we may assume that κ -compact objects are stable under R -limits.*
- *We may assume that many such properties hold for the same cardinal.*

Definition

A morphism $f : X \rightarrow Y$ in an ∞ -category is said to be relatively κ -compact if for every κ -compact object Z and every diagram

$$\begin{array}{ccc} W & \longrightarrow & X \\ \downarrow & \lrcorner & \downarrow f \\ Z & \longrightarrow & Y \end{array}$$

the object W is also κ -compact.

Definition

A morphism $f : X \rightarrow Y$ in an ∞ -category is said to be relatively κ -compact if for every κ -compact object Z and every diagram

$$\begin{array}{ccc} W & \longrightarrow & X \\ \downarrow & \lrcorner & \downarrow f \\ Z & \longrightarrow & Y \end{array}$$

the object W is also κ -compact.

Theorem (Rezk)

In a geometric ∞ -topos, there are arbitrarily large cardinals κ such that the class S_κ of relatively κ -compact morphisms has a classifier.

Theorem

Fixing a Grothendieck universe \mathcal{U} , every geometric ∞ -topos satisfies (DepProd) if and only if there are unboundedly many inaccessible cardinals below the cardinality of \mathcal{U} .

Theorem

Fixing a Grothendieck universe \mathcal{U} , every geometric ∞ -topos satisfies (DepProd) if and only if there are unboundedly many inaccessible cardinals below the cardinality of \mathcal{U} .

First, prove \Leftarrow .

We want to use Rezk's theorem to find universes in the form S_{κ} . We will need uniformization and the hypothesis to find suitable κ 's.

Theorem

Fixing a Grothendieck universe \mathcal{U} , every geometric ∞ -topos satisfies (DepProd) if and only if there are unboundedly many inaccessible cardinals below the cardinality of \mathcal{U} .

First, prove \Leftarrow .

We want to use Rezk's theorem to find universes in the form S_{κ} . We will need uniformization and the hypothesis to find suitable κ 's.

Step 1. In the ∞ -category \mathcal{S} of spaces, if κ is inaccessible then κ -compact objects are stable under exponentiation.

Step 2. In $\mathcal{P}(\mathcal{C})$, given presheaves F and G , their exponential F^G is given by the formula

$$F^G(C) = \int_{D \in \mathcal{C}} \text{Map}(\text{Map}(D, C) \times G(D), F(D))$$

Step 2. In $\mathcal{P}(\mathcal{C})$, given presheaves F and G , their exponential F^G is given by the formula

$$F^G(C) = \int_{D \in \mathcal{C}} \text{Map}(\text{Map}(D, C) \times G(D), F(D))$$

By uniformization, we may choose a cardinal κ such that:

- κ -compactness is detected objectwise

Step 2. In $\mathcal{P}(\mathcal{C})$, given presheaves F and G , their exponential F^G is given by the formula

$$F^G(C) = \int_{D \in \mathcal{C}} \text{Map}(\text{Map}(D, C) \times G(D), F(D))$$

By uniformization, we may choose a cardinal κ such that:

- κ -compactness is detected objectwise
- all representables are κ -compact

Step 2. In $\mathcal{P}(\mathcal{C})$, given presheaves F and G , their exponential F^G is given by the formula

$$F^G(C) = \int_{D \in \mathcal{C}} \text{Map}(\text{Map}(D, C) \times G(D), F(D))$$

By uniformization, we may choose a cardinal κ such that:

- κ -compactness is detected objectwise
- all representables are κ -compact
- κ -compact spaces are stable under binary products

Step 2. In $\mathcal{P}(\mathcal{C})$, given presheaves F and G , their exponential F^G is given by the formula

$$F^G(C) = \int_{D \in \mathcal{C}} \text{Map}(\text{Map}(D, C) \times G(D), F(D))$$

By uniformization, we may choose a cardinal κ such that:

- κ -compactness is detected objectwise
- all representables are κ -compact
- κ -compact spaces are stable under binary products
- κ -compact spaces are stable under exponentiation (Step 1)

Step 2. In $\mathcal{P}(\mathcal{C})$, given presheaves F and G , their exponential F^G is given by the formula

$$F^G(C) = \int_{D \in \mathcal{C}} \text{Map}(\text{Map}(D, C) \times G(D), F(D))$$

By uniformization, we may choose a cardinal κ such that:

- κ -compactness is detected objectwise
- all representables are κ -compact
- κ -compact spaces are stable under binary products
- κ -compact spaces are stable under exponentiation (Step 1)
- κ -compact spaces are stable under \mathcal{C} -ends

Step 2. In $\mathcal{P}(\mathcal{C})$, given presheaves F and G , their exponential F^G is given by the formula

$$F^G(C) = \int_{D \in \mathcal{C}} \text{Map}(\text{Map}(D, C) \times G(D), F(D))$$

By uniformization, we may choose a cardinal κ such that:

- κ -compactness is detected objectwise
- all representables are κ -compact
- κ -compact spaces are stable under binary products
- κ -compact spaces are stable under exponentiation (Step 1)
- κ -compact spaces are stable under \mathcal{C} -ends

\Rightarrow κ -compact presheaves are stable under exponentiation.

Step 3. Given an adjunction

$$\mathcal{P}(\mathcal{C}) \begin{array}{c} \xrightarrow{L} \\ \xleftarrow[\perp]{i} \end{array} \mathcal{X}$$

making \mathcal{X} a geometric ∞ -topos, choose κ such that (Step 2) holds in $\mathcal{P}(\mathcal{C})$.

Step 3. Given an adjunction

$$\mathcal{P}(\mathcal{C}) \begin{array}{c} \xrightarrow{L} \\ \xleftarrow{i} \\ \perp \end{array} \mathcal{X}$$

making \mathcal{X} a geometric ∞ -topos, choose κ such that (Step 2) holds in $\mathcal{P}(\mathcal{C})$.

The properties of $L \dashv i$ will transfer stability of κ -compact objects under exponentiation to \mathcal{X} .

Step 4. Given an object $p : Z \rightarrow X$ in \mathcal{X}/X , its dependent product along a terminal morphism $X \rightarrow *$ is given by

$$\prod_X p = Z^X \times_{X^X} \{p\}$$

Step 4. Given an object $p : Z \rightarrow X$ in \mathcal{X}/X , its dependent product along a terminal morphism $X \rightarrow *$ is given by

$$\prod_X p = Z^X \times_{X^X} \{p\}$$

Choose κ such that (Step 3) holds and κ -compact objects are stable under pullbacks

Step 4. Given an object $p : Z \rightarrow X$ in \mathcal{X}/X , its dependent product along a terminal morphism $X \rightarrow *$ is given by

$$\prod_X p = Z^X \times_{X^X} \{p\}$$

Choose κ such that (Step 3) holds and κ -compact objects are stable under pullbacks \Rightarrow relatively κ -compact morphisms are stable under dependent products along terminal morphisms.

Step 4. Given an object $p : Z \rightarrow X$ in \mathcal{X}/X , its dependent product along a terminal morphism $X \rightarrow *$ is given by

$$\prod_X p = Z^X \times_{X^X} \{p\}$$

Choose κ such that (Step 3) holds and κ -compact objects are stable under pullbacks \Rightarrow relatively κ -compact morphisms are stable under dependent products along terminal morphisms.

Step 5. For generic dependent products, decompose the codomain as a colimit of compact objects Y_i 's and then choose κ such that (Step 4) holds in all ∞ -toposes \mathcal{X}/Y_i .

Main result

Now prove \Rightarrow .

Main result

Now prove \Rightarrow . It suffices to prove it assuming that \mathcal{S} satisfies $(DepProd)$.

Main result

Now prove \Rightarrow . It suffices to prove it assuming that \mathcal{S} satisfies $(DepProd)$.

For a discrete space X , the terminal morphism $X \rightarrow *$ is contained in a class \mathcal{S} having a classifier $t : \bar{U} \rightarrow U$ such that

$$\begin{array}{ccc} Y & \longrightarrow & \bar{U} \\ p \downarrow & \lrcorner & t \downarrow \\ Z & \longrightarrow & U \end{array}, \quad \begin{array}{ccc} Z & \longrightarrow & \bar{U} \\ f \downarrow & \lrcorner & t \downarrow \\ W & \longrightarrow & U \end{array}$$

Main result

Now prove \Rightarrow . It suffices to prove it assuming that \mathcal{S} satisfies $(DepProd)$.

For a discrete space X , the terminal morphism $X \rightarrow *$ is contained in a class \mathcal{S} having a classifier $t : \bar{U} \rightarrow U$ such that

$$\begin{array}{ccc} Y \longrightarrow \bar{U} & & Z \longrightarrow \bar{U} \\ p \downarrow \lrcorner & & f \downarrow \lrcorner \\ Z \longrightarrow U & , & W \longrightarrow U \end{array} \quad \Longrightarrow \quad \exists \begin{array}{ccc} \prod_f p \longrightarrow \bar{U} & & \\ \downarrow \lrcorner & & t \downarrow \\ W \longrightarrow U & & \end{array}$$

Main result

Now prove \Rightarrow . It suffices to prove it assuming that \mathcal{S} satisfies $(DepProd)$.

For a discrete space X , the terminal morphism $X \rightarrow *$ is contained in a class \mathcal{S} having a classifier $t : \bar{U} \rightarrow U$ such that

$$\begin{array}{ccc} Y \longrightarrow \bar{U} & & Z \longrightarrow \bar{U} \\ p \downarrow \lrcorner & & f \downarrow \lrcorner \\ Z \longrightarrow U & & W \longrightarrow U \end{array}, \quad \Rightarrow \quad \exists \quad \begin{array}{ccc} \prod_f p \longrightarrow \bar{U} & & \\ \downarrow \lrcorner & & t \downarrow \\ W \longrightarrow U & & \end{array}$$

- Assume that all fibers of t are discrete.

Main result

Now prove \Rightarrow . It suffices to prove it assuming that \mathcal{S} satisfies $(DepProd)$.

For a discrete space X , the terminal morphism $X \rightarrow *$ is contained in a class \mathcal{S} having a classifier $t : \bar{U} \rightarrow U$ such that

$$\begin{array}{ccc} Y \longrightarrow \bar{U} & & Z \longrightarrow \bar{U} \\ p \downarrow \lrcorner & & f \downarrow \lrcorner \\ Z \longrightarrow U & , & W \longrightarrow U \end{array} \quad \Longrightarrow \quad \exists \begin{array}{ccc} \prod_f p \longrightarrow \bar{U} & & \\ \downarrow \lrcorner & & t \downarrow \\ W \longrightarrow U & & \end{array}$$

- Assume that all fibers of t are discrete.
- For each point in U , its fiber along t can be regarded as a set.

Main result

$$\begin{array}{ccc} F_x & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x\} & \longrightarrow & U \end{array}$$

$$\begin{array}{ccc} X & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x_0\} & \longrightarrow & U \end{array}$$

Main result

$$\begin{array}{ccc} F_x & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x\} & \longrightarrow & U \end{array}$$

$$\begin{array}{ccc} X & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x_0\} & \longrightarrow & U \end{array}$$

Define $\kappa := \sup_{x \in U} |F_x|$.

Main result

$$\begin{array}{ccc} F_x & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x\} & \longrightarrow & U \end{array}$$

$$\begin{array}{ccc} X & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x_0\} & \longrightarrow & U \end{array}$$

Define $\kappa := \sup_{x \in U} |F_x|$.

- $\kappa > |X|$.

Main result

$$\begin{array}{ccc} F_x & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x\} & \longrightarrow & U \end{array}$$

$$\begin{array}{ccc} X & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x_0\} & \longrightarrow & U \end{array}$$

Define $\kappa := \sup_{x \in U} |F_x|$.

- $\kappa > |X|$.
- For $\lambda, \mu < \kappa$, closure under dependent products $\Rightarrow \mu^\lambda < \kappa$.

Main result

$$\begin{array}{ccc} F_x & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x\} & \longrightarrow & U \end{array}$$

$$\begin{array}{ccc} X & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x_0\} & \longrightarrow & U \end{array}$$

Define $\kappa := \sup_{x \in U} |F_x|$.

- $\kappa > |X|$.
- For $\lambda, \mu < \kappa$, closure under dependent products $\Rightarrow \mu^\lambda < \kappa$.
- In non-trivial cases, $\sum_{i \in I} \alpha_i \leq \prod_{i \in I} \alpha_i$

Main result

$$\begin{array}{ccc} F_x & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x\} & \longrightarrow & U \end{array}$$

$$\begin{array}{ccc} X & \longrightarrow & \bar{U} \\ \downarrow & \lrcorner & \downarrow t \\ \{x_0\} & \longrightarrow & U \end{array}$$

Define $\kappa := \sup_{x \in U} |F_x|$.

- $\kappa > |X|$.
- For $\lambda, \mu < \kappa$, closure under dependent products $\Rightarrow \mu^\lambda < \kappa$.
- In non-trivial cases, $\sum_{i \in I} \alpha_i \leq \prod_{i \in I} \alpha_i \Rightarrow \kappa$ is regular.

Definition

We call a cardinal μ 1-inaccessible if it is inaccessible and there are unboundedly many inaccessibles below it.

Definition

We call a cardinal μ 1-inaccessible if it is inaccessible and there are unboundedly many inaccessibles below it.

Assume the existence of a 1-inaccessible cardinal μ inside the Grothendieck universe.

Definition

We call a cardinal μ 1-inaccessible if it is inaccessible and there are unboundedly many inaccessibles below it.

Assume the existence of a 1-inaccessible cardinal μ inside the Grothendieck universe.

Given a geometric ∞ -topos \mathcal{X} , take

$$\mathcal{X}^\mu \subset \mathcal{X}.$$

Definition

We call a cardinal μ 1-inaccessible if it is inaccessible and there are unboundedly many inaccessibles below it.

Assume the existence of a 1-inaccessible cardinal μ inside the Grothendieck universe.

Given a geometric ∞ -topos \mathcal{X} , take

$$\mathcal{X}^\mu \subset \mathcal{X}.$$

$\Rightarrow \mathcal{X}^\mu$ is not a geometric ∞ -topos (it doesn't have all small colimits), but it is an elementary ∞ -topos.

Thank you!