

# On epimorphisms and acyclic types in $\text{HoTT}^*$

Ulrik Buchholtz<sup>1</sup>, Tom de Jong<sup>1</sup>, and Egbert Rijke<sup>2</sup>

<sup>1</sup> School of Computer Science, University of Nottingham

<sup>2</sup> Department of Mathematics, University of Ljubljana

We study epimorphisms and acyclic types in univalent mathematics. The epimorphisms are of course a natural object of study, the definition being that a map  $f : A \rightarrow B$  is an *epimorphism* if for every type  $X$ , the precomposition map

$$(B \rightarrow X) \xrightarrow{(-) \circ f} (A \rightarrow X) \quad (1)$$

is an embedding. In other words, extensions of maps out of  $A$  through  $B$  are unique when they exist.

The epimorphisms behave quite differently in higher types than do the surjections of sets, with the difference manifesting itself starting at the level of 2-types. To bring out this structure, we introduce the notion of *k-epimorphism*, which is a map  $f$  such that (1) is an embedding for all  $k$ -types  $X$ .

We prove that being a ( $k$ -)epimorphism is a fiberwise condition. The 0-epimorphisms are the surjections, i.e., the  $(-1)$ -connected maps, and the 1-epimorphisms are the 0-connected maps. But the 2-epimorphisms are the maps whose fibers are connected and whose fundamental groups are *perfect*. (A group  $G$  is perfect if its abelianization is trivial.)

This connects the notion of epimorphism with that of *acyclic* types. These are classically the types  $A$  whose reduced integral homology vanishes,  $\tilde{H}_n(A) \simeq 0$  for all  $n \geq 0$ . This characterization only works assuming Whitehead's principle, and the correct definition in univalent mathematics is that  $A$  is acyclic if its suspension,  $\Sigma A$ , is contractible. We prove:

**Proposition 1.** *A map  $A \rightarrow \mathbf{1}$  is  $k$ -epic if and only if  $\Sigma A$  is  $k$ -connected, and an epimorphism if and only if  $\Sigma A$  is contractible.*

Correspondingly, we call a type  $A$  *k-acyclic* if  $\Sigma A$  is  $k$ -connected.

For a group  $G$  with classifying type  $BG$ , we know that the abelianization  $G^{\text{ab}}$  has classifying type  $B^2G^{\text{ab}} \simeq \|\Sigma BG\|_2$  [BvDR18]. Thus we obtain:

**Corollary 2.** *For a group  $G$ , the map  $BG \rightarrow \mathbf{1}$  is 2-epic if and only if  $G$  is perfect.*

Next we begin the search for examples of acyclic types. These cannot be sets because of the following:

**Theorem 3.** *For a set  $A$ , the following are equivalent:*

- (i)  $A$  is 1-acyclic;
- (ii)  $A$  is acyclic;
- (iii)  $A$  is contractible.

---

\*This abstract was also submitted to the *Workshop on Homotopy Type Theory/Univalent Foundations* which will take place on 22 and 23 April in Vienna, Austria.

Interestingly, to prove Theorem 3, we need the fact that the generators map  $\eta : A \rightarrow F_A$ , from  $A$  to the free group on  $A$  is an injection [MRR88, Chapter X]. This has previously been proved in homotopy type theory [BCDE21].

We also note the following, whose proof is short enough to fit here:

**Lemma 4.** *Any acyclic and simply connected type  $A$  is infinitely connected.*

*Proof.* This follows by the Freudenthal suspension theorem: The unit map of the loop-suspension adjunction,  $\sigma : A \rightarrow \Omega\Sigma A$ , is  $2n$ -connected whenever  $A$  is  $n$ -connected (for  $n \geq 0$ ). By acyclicity,  $\Sigma A = \mathbf{1}$ , so  $\Omega\Sigma A = \mathbf{1}$ , so the connectivity of  $\sigma$  is that of  $A$ . Starting with the assumption that  $A$  is 1-connected, we in turn conclude that  $A$  is 2-, then 4-, etc., hence  $2^n$ -connected, for all  $n$ .  $\square$

Classically, it's well known that the Higman group [Hig51], given by the presentation

$$H = \langle a, b, c, d \mid a = [d, a], b = [a, b], c = [b, c], d = [c, d] \rangle$$

is acyclic (i.e., has an acyclic classifying space), and moreover, this presentation is *aspheric*, meaning that the presentation complex is already a 1-type [DV73]. The presentation complex is easily imported into homotopy type theory as the higher inductive type  $BH$  with a point constructor  $\text{pt} : BH$ , four path constructors  $a, b, c, d : \Omega BH$ , and four 2-cell constructors corresponding to the relations. We prove:

**Proposition 5.** *The type  $BH$  is acyclic and  $\pi_1(BH)$  is the Higman group.*

We leave for future work the matter of developing enough combinatorial group theory to show that  $BH$  is a 1-type.

We note that nullification at  $BH$  provides a nontrivial modality  $L$ , whose subuniverse of  $L$ -separated types consists of all types, following [CR22, Ex. 6.6].

**Future directions** Classically, the acyclic maps form the left class of an accessible orthogonal factorization system, whose right class are the *hypoabelian* maps (i.e., whose  $\pi_1$ -kernels have no non-trivial perfect subgroups). This factorization system exists in all  $(\infty, 1)$ -toposes [Hoy19], and can be used to derive the McDuff–Segal completion theorem, but with arguments that don't lend themselves to direct internalization in homotopy type theory.

We only prove the main closure properties of the acyclic maps as one should expect from such a left class. We hope in future work to construct the factorization system using Quillen's *plus construction*. To do so, we expect to need the strengthening of Lemma 4 stating that acyclic, simply connected types are actually contractible. It appears to be an open problem whether this holds in all  $(\infty, 1)$ -toposes. It follows from Whitehead's principle, but also holds in  $(\infty, 1)$ -toposes where this fails, such as that of parametrized spectra. So we propose to assume it as a principle, which we might call the *plus principle*.

We would also like to investigate whether other classically known acyclic types and maps can be shown to be so in homotopy type theory, such as  $B \text{Aut}(\mathbb{N})$  [dlHM83] and  $B\Sigma_\infty \rightarrow (\Omega^\infty \Sigma^\infty \mathbb{S}^0)_0$ , the Barratt–Priddy(–Quillen) theorem [BP72].

## References

- [BCDE21] Marc Bezem, Thierry Coquand, Peter Dybjer and Martín Escardó. ‘Free groups in HoTT/UF in Agda’. <https://www.cs.bham.ac.uk/~mhe/TypeTopology/FreeGroup.html>. 2021.

- [BP72] Michael Barratt and Stewart Priddy. ‘On the homology of non-connected monoids and their associated groups’. In: *Comment. Math. Helv.* 47 (1972), pp. 1–14. DOI: [10.1007/BF02566785](https://doi.org/10.1007/BF02566785).
- [BvDR18] Ulrik Buchholtz, Floris van Doorn and Egbert Rijke. ‘Higher Groups in Homotopy Type Theory’. In: *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science. LICS ’18*. Oxford, United Kingdom: Association for Computing Machinery, 2018, pp. 205–214. DOI: [10.1145/3209108.3209150](https://doi.org/10.1145/3209108.3209150). arXiv: [1802.04315](https://arxiv.org/abs/1802.04315).
- [CR22] J. Daniel Christensen and Egbert Rijke. ‘Characterizations of modalities and lex modalities’. In: *J. Pure Appl. Algebra* 226.3 (2022), Paper No. 106848, 21. DOI: [10.1016/j.jpaa.2021.106848](https://doi.org/10.1016/j.jpaa.2021.106848).
- [dlHM83] Pierre de la Harpe and Dusa McDuff. ‘Acyclic groups of automorphisms’. In: *Comment. Math. Helv.* 58.1 (1983), pp. 48–71. DOI: [10.1007/BF02564624](https://doi.org/10.1007/BF02564624).
- [DV73] Eldon Dyer and A. T. Vasquez. ‘Some small aspherical spaces’. In: *Journal of the Australian Mathematical Society* 16.3 (1973), pp. 332–352. DOI: [10.1017/S1446788700015147](https://doi.org/10.1017/S1446788700015147).
- [Hig51] Graham Higman. ‘A finitely generated infinite simple group’. In: *J. London Math. Soc.* 26 (1951), pp. 61–64. DOI: [10.1112/jlms/s1-26.1.61](https://doi.org/10.1112/jlms/s1-26.1.61).
- [Hoy19] Marc Hoyois. ‘On Quillen’s plus construction’. 2019. URL: <https://hoyois.app.uni-regensburg.de/papers/acyclic.pdf>.
- [MRR88] Ray Mines, Fred Richman and Wim Ruitenburg. *A Course in Constructive Algebra*. Universitext. Springer, 1988. DOI: [10.1007/978-1-4419-8640-5](https://doi.org/10.1007/978-1-4419-8640-5).